PROCEEDINGS OF SCIENCE

arXiv:1511.00477v1 [hep-lat] 2 Nov 2015

Large N meson propagators from twisted space-time reduced model

Antonio González-Arroyo[ab]
[a]*Instituto de Física Teórica UAM/CSIC, C/ Nicolás Cabrera 13-15*
Universidad Autónoma de Madrid, E-28048–Madrid, Spain
[b]*Departamento de Física Teórica, C-15*
Universidad Autónoma de Madrid, E-28049–Madrid, Spain
E-mail: antonio.gonzalez-arroyo@uam.es

Masanori Okawa[*cd]
[c]*Graduate School of Science, Hiroshima University*
Higashi-Hiroshima, Hiroshima 739-8526, Japan
[d]*Core of Research for the Energetic Universe, Hiroshima University*
Higashi-Hiroshima, Hiroshima 739-8526, Japan
E-mail: okawa@sci.hiroshima-u.ac.jp

Recently, we proposed a new method to calculate meson propagators in the large N limit from twisted space-time reduced model. In this note, we give simulation details for obtaining meson spectra and discuss the smearing technique which should improve the signal of meson propagators in future works.

The 33rd International Symposium on Lattice Field Theory
14 -18 July 2015
*Kobe International Conference Center, Kobe, Japan**

[*]Speaker.

© Copyright owned by the author(s) under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND 4.0).

1. Introduction

In the last few years, we have performed several studies of the twisted space-time reduced model. This is a one-site lattice model[1, 2], expected to become equivalent in the large N limit to the usual lattice gauge theory at infinite volume. We have identified the condition on the twisted boundary condition which ensures this equivalence[3]. We have also calculated a variety of quantities on both lattice[4] and continuum[5, 6] space-time. We also showed that it is straightforward to introduce adjoint fermions within our framework[7, 8].

Although one of the most important observables of lattice gauge theories is given by the hadron spectra, our previous studies did not attempt to calculate it in the twisted reduced model. At first sight it seemed impossible to calculate space-time extended hadron correlation functions in the one-site model. Furthermore, the presence of quarks in the fundamental representation of the group seemed in conflict with twisted boundary conditions. Recently, we found a way to circumvent these difficulties and succeeded in calculating meson spectra in the large N limit[9]. The purpose of this note is to present simulation details which are not covered in ref[9], and propose a new smearing method which should be indispensable for a precise determination of meson spectra.

2. Formulation

We consider the twisted Eguchi-Kawai model with gauge group SU(N) with $N = L^2$, L being positive (preferably prime) integer. The action of the TEK model is given by

$$S = -bN \sum_{\mu \neq \nu = 0}^{d-1} z_{\nu\mu} \mathrm{Tr}\left[U_\mu U_\nu U_\mu^\dagger U_\nu^\dagger\right] \tag{2.1}$$

Here, U_μ are d=4 SU(N) link variables. The symmetric twist tensor $z_{\mu\nu}$ is an element of Z(L);

$$z_{\nu\mu} = \exp\left(k\frac{2\pi i}{L}\right), \quad z_{\mu\nu} = z_{\nu\mu}^*, \quad \mu > \nu \tag{2.2}$$

k and L are co-prime and should obey some constraint in order to avoid Z(N) symmetry breaking at intermediate values of the coupling[3]. The minimum action solution is given by matrices Γ_μ satisfying

$$\Gamma_\mu \Gamma_\nu = z_{\mu\nu} \Gamma_\nu \Gamma_\mu \tag{2.3}$$

whose explicit form is given, for example, in eq. (2.13) of ref.[4].

As has been shown in ref.[9], the meson propagator in channel γ_A and γ_B at time distance n_0 is given by the following formula

$$C_{AB}(n_0) = \frac{1}{\ell_0 N^{3/2}} \sum_{q_0} \mathrm{e}^{-iq_0 n_0} \mathrm{Tr}\left[\gamma_A D^{-1}(0) \gamma_B D^{-1}(q_0)\right] \tag{2.4}$$

Here $D(q_0)$ is the Wilson-Dirac matrix acting on color (U_μ), spatial (Γ_μ) and Dirac (γ_μ) spaces and its explicit form is

$$D(q_0) = 1 - \kappa \sum_{\mu=0}^{d-1} \left[(1-\gamma_\mu)\tilde{U}_\mu \Gamma_\mu^* + (1+\gamma_\mu)\tilde{U}_\mu^\dagger \Gamma_\mu^t\right] \tag{2.5}$$

with

$$\tilde{U}_{\mu=0} = \mathrm{e}^{iq_0} U_{\mu=0}, \quad \tilde{U}_{\mu=1,2,3} = U_{\mu=1,2,3} \tag{2.6}$$

and

$$q_0 = \frac{2\pi m}{\ell_0 L}, \quad 0 \le m \le \ell_0 L - 1 \tag{2.7}$$

Quark fields are supposed to live in a finite box $\ell_0 L \times L^3$ with positive integer ℓ_0, whereas gauge fields live in a $L \times L^3$. A value of $\ell_0 > 1$ is very convenient since in formula (2.4) we represent the correlation function of ultralocal meson operators in spatially zero momentum state. This correlator receives important contributions of higher excitation and one must go to relatively long time separation to extract a good signal for the lowest mass state. As will be discussed in sect. 4, this problem can be circumvented by using spatially extended smeared meson operators.

We use $Z(4)$ random source method to compute Tr in (2.4). Let $z(\alpha,\beta,i)$ be the source vector having color (U_μ) index $1 \le \alpha \le N$, spatial (Γ_μ) index $1 \le \beta \le N$ and Dirac (γ_μ) index $1 \le i \le 4$. Then $z(\alpha,\beta,i) = \frac{1}{\sqrt{2}}(\pm 1 \pm i)$. After averaging over random source, we have $\langle z^*(\alpha',\beta',i') z(\alpha,\beta,i)\rangle = \delta_{\alpha'\alpha}\delta_{\beta'\beta}\delta_{i'i}$. Now we define Hermitian operator $Q(q_0) = D(q_0)\gamma_5$ and solve the following $\ell_0 L + 1$ equations.

$$Q(0) x_A = \gamma_5 \gamma_A^\dagger z \tag{2.8}$$

$$Q(q_0)\gamma_5\gamma_B^{-1} y_B(q_0) = z, \quad q_0 = \frac{2\pi m}{\ell_0 L}, \quad 0 \le m \le \ell_0 L - 1 \tag{2.9}$$

Taking the inner product of $x_A^\dagger$ and $y_B(q_0)$ and averaging over random source, we have

$$\begin{aligned} \left\langle x_A^\dagger y_B(q_0) \right\rangle &= \left\langle z^\dagger \gamma_A \gamma_5 Q^{-1}(0) \gamma_B \gamma_5 Q^{-1}(q_0) z \right\rangle \\ &= \mathrm{Tr}\left[\gamma_A D^{-1}(0) \gamma_B D^{-1}(q_0)\right] \end{aligned} \tag{2.10}$$

We use the CG inversion, so Q^{-1} actually means $Q^{-1} = QQ^{-2}$. It should be noted that once we have calculated $y_5(q_0)$ for all q_0, $y_B(q_0)$ can be obtained without the CG inversion as

$$y_B(q_0) = \gamma_B \gamma_5 y_5(q_0) \tag{2.11}$$

3. Simulation

The $d = 4$ gauge link variables U_μ are generated by a recently proposed over-relaxation Monte Carlo method[10]. This method produces independent gauge configuration approximately twice faster than the conventional heat bath method[11]. We choose $N = 289 = 17^2$. Therefore the effective lattice size of the gauge field is 17^4. We take $k = 5$ to ensure that our theory does not suffer from Z(N) symmetry breaking and use 5 random sources to approximate (2.10). We study two values of gauge coupling $b = 0.36$ and 0.37, and for each b, 800 gauge configurations are stored. Each configuration are separated by 1000 MC sweeps. Errors are estimated by jackknife method.

In fig. 1, we show $\frac{1}{N}\mathrm{Tr}\left[\gamma_5 D^{-1}(0)\gamma_5 D^{-1}(q_0)\right]$ with $q_0 = \frac{2\pi m}{\ell_0 L}$ for $\ell_0 = 2$, at b=0.36 and κ=0.155. The actual error bars are too small to see in this scale, so they are artificially enhanced 100 times. We notice error bars are larger for smaller q_0 (mod 2π).



Figure 1: $\frac{1}{N}\mathrm{Tr}\left[\gamma_5 D^{-1}(0)\gamma_5 D^{-1}(q_0=\frac{2\pi m}{\ell_0 L})\right]$ as a function of m at b=0.36 and κ=0.155. The error bars of y-axis are 100 times enhanced artificially.

Meson correlation functions are extracted from (2.4). By making the sum over $q_0 = \frac{2\pi m}{\ell_0 L}$, $0 \le m \le \ell_0 L - 1$ for $0 \le n_0 \le \ell_0 L - 1$, we obtain the correlation function for the quarks living in a $\ell_0 L \times L^3$ finite box.

In fig. 2, we show the pion correlation function $C_{55}(n_0)$ both for $\ell_0 = 1$ and 2. For ℓ_0 =1, signals are good for entire range of $0 \le n_0 \le L = 17$. However, contaminations from higher excited states are so large that we can not reliably extract the information of the lowest state. On the other hand, for ℓ_0 =2, although the data points are quite smooth as a function of n_0, the error bars become large for larger separation. This is in sharp contrast to the convention calculation of pion correlator, where we know that the error of the pion correrator decreases proportional to the value of the correlator itself. The reason is simple; we extract the correlation function from (2.4). As is pointed out earler, largest error comes from $\mathrm{Tr}\left[\gamma_5 D^{-1}(0)\gamma_5 D^{-1}(0)\right]$, whose contribution to $C_{AB}(n_0)$ is independent on n_0.

We, therefore, expect that constant error contributions are canceled out in effective mass $m_{\mathrm{eff}}(n_0) = \log[C(n_0-1)/C(n_0)]$ and get good signals even for $\ell_0 = 2$. Results are shown in fig. 3 for both pion (black) and rho meson(red) at b=0.36 and κ=0.155. We clearly see good signals up to $n_0 \sim 13$, although contributions of higher excitation states are not negligible for small n_0 due to the use of local meson operators. We then make a three mass fit of the form

$$m_{\mathrm{eff}}(n_0) = \log\left[G(n_0-1)/G(n_0)\right] \tag{3.1}$$

$$G(n_0) = \mathrm{e}^{-m_1 n_0} + a_2 \mathrm{e}^{-m_2 n_0} + a_3 \mathrm{e}^{-m_3 n_0} \tag{3.2}$$

with the fitting range $[3:13]$ for pion and $[2:13]$ for rho meson. The fits are reasonable with χ^2/ndf 0.76 for pion and 0.63 for rho meson. We have repeated the above procedure for other κ values with b=0.36 and 0.37, and these results are used to make discussion in ref.[9].



Figure 2: Pion correlation function $C_{55}(n_0)$ as functions of n_0 at b=0.36 and κ=0.155. Green symbols represent the results with $\ell_0 = 1$ whereas red symbols are for $\ell_0 = 2$.



Figure 3: Effective mass $m_{\rm eff}(n_0) = \log[C(n_0-1)/C(n_0)]$ as functions of n_0 for pion (black) and rho meson(red). Fits are made with fitting function (3.1).

4. Smearing

From the above discussion, it is obvious that we have to reduce the effects of higher excitation states to get more reliable results. Usually, this is achieved by constructing spatially extended operators having the same quantum numbers and make a variational analysis. At first sight, it does not seem to be clear how to make spatially extended operators in our one-site model. Nevertheless, it is a rather straightforward generalization of our formulation to construct smeared operators.

Our proposal is summarized as follow. For the meson operator in channel A, we replace γ_A by



Figure 4: Effective mass $m_{\rm eff}^{ll}(n_0) = \log\left[C_{55}^{ll}(n_0-1)/C_{55}^{ll}(n_0)\right]$ for pion as functions of n_0 in $d=2$ TEK model. Red symbols are for unsmeared operator and green symbols are 15 times smeared operator.

the following operator having the same quantum number

$$\gamma_A \to D_s^l \gamma_A, \quad D_s \equiv \frac{1}{1+6c}\left[1 + c\sum_{i=1}^{d-1}\left(\bar{U}_i\Gamma_i^* + \bar{U}_i^\dagger\Gamma_i^t\right)\right] \tag{4.1}$$

l is the smearing level and $\bar{U}_i$ is the ape-smeared spatial link variables after making the following transformation several times iteratively

$$U_i' = \mathrm{Proj}_{\mathrm{SU(N)}}\left[(1-f)U_i + \frac{f}{4}\sum_{j\neq i}(U_jU_iU_j^\dagger + U_j^\dagger U_iU_j)\right] \tag{4.2}$$

c and f are smearing parameters. (4.1) is just the one-site version of the smearing adopted in [12]. The smeared meson propagators are given by

$$C_{AB}^{ll'}(n_0) = \frac{1}{\ell_0 N^{3/2}}\sum_{q_0} e^{-iq_0n_0}\mathrm{Tr}\left[D_s^l\gamma_A D^{-1}(0) D_s^{l'}\gamma_B D^{-1}(q_0)\right], \tag{4.3}$$

which can be easily calculated from

$$x_A^l = Q^{-1}(0)D_s^l\gamma_5\gamma_A^\dagger z \tag{4.4}$$

$$y_B^{l'}(q_0) = D_s^{l'}y_B(q_0) \tag{4.5}$$

To check the validity of the smearing, we have applied it to the $d=2$ TEK model. In two dimension, SU(N) TEK model is related to the usual lattice theory on a $N\times N$ finite box. Since there is no extra spatial dimension to make ape-smearing, D_s takes very simple form

$$D_s = \frac{1}{1+2c}\left[1 + c\left(U_1\Gamma_1^* + U_1^\dagger\Gamma_1^t\right)\right] \tag{4.6}$$

Simulations are made with N=31, k=7, b=2.0 and κ=0.25.

In fig. 4, we show the effective mass of pion both for unsmeared local operator ($l = 0$, red symbol) and that of 15 times smeared spatially extended operator ($l = 15$, $c = 0.4$, green symbol) with $\ell_0 = 1$. The effect of the smearing is significant. We are now making systematic variational analysis for both 2 and 4 dimensional TEK model[13]. We hope to present new results in coming lattice conference.

Acknowledgments

We acknowledge financial support from the MCINN grants FPA2012-31686 and FPA2012-31880, and the Spanish MINECO's "Centro de Excelencia Severo Ochoa" Programme under grant SEV-2012-0249. M. O. is supported by the Japanese MEXT grant No 26400249 and the MEXT program for promoting the enhancement of research universities. Calculations have been done on Hitachi SR16000 supercomputer both at High Energy Accelerator Research Organization(KEK) and YITP in Kyoto University. Work at KEK is supported by the Large Scale Simulation Program No.15/16-04.

References

[1] A. Gonzalez-Arroyo and M. Okawa, Phys. Lett. B **120** (1983) 174.

[2] A. Gonzalez-Arroyo and M. Okawa, Phys. Rev. D **27**, 2397 (1983).

[3] A. Gonzalez-Arroyo and M. Okawa, JHEP **07** (2010) 043 [arXiv:1005.1981 [hep-th]].

[4] A. Gonzalez-Arroyo and M. Okawa, JHEP **12** (2014) 106 [arXiv:1410.6405 [hep-lat]].

[5] A. Gonzalez-Arroyo and M. Okawa, Phys. Lett. B **718** (2013) 1524 [arXiv:1206.0049 [hep-th]].

[6] M. Garcia Perez, A. Gonzalez-Arroyo, L. Keegan and M. Okawa, JHEP **01** (2015) 038 [arXiv:1412.0941 [hep-lat]].

[7] A. Gonzalez-Arroyo and M. Okawa, Phys. Rev. D **88** (2013) 014514 [arXiv:1305.6253 [hep-lat]].

[8] M. Garcia Perez, A. Gonzalez-Arroyo, L. Keegan and M. Okawa, JHEP **08** (2015) 034 [arXiv:1506.06536 [hep-lat]].

[9] A. Gonzalez-Arroyo and M. Okawa, [arXiv:1510.05428 [hep-lat]].

[10] M. Garcia Perez, A. Gonzalez-Arroyo, L. Keegan, M. Okawa and A. Ramos, JHEP **06** (2015) 093 [arXiv:1505.05784 [hep-lat]].

[11] K. Fabricius and O. Haan, Phys. Lett. B **143** (1984) 459.

[12] G. Bali, F. Bursa, L. Castagnini, S. Collins, L. Del Debbio, B. Lucini and M. Panero, JHEP **06** (2013) 071.

[13] M. Garcia Perez, A. Gonzalez-Arroyo and M. Okawa, in preparation.